


# Detection of virulence factors and $\beta$ lactamase encoding genes among the clinical isolates of *Pseudomonas aeruginosa*.

Fazlul MKK<sup>1</sup>, Najnin A<sup>2</sup>, Farzana Y<sup>3</sup>, Rashid MA<sup>4</sup>, Deepthi S<sup>5</sup>, Srikumar C<sup>5</sup>, SS Rashid<sup>1</sup>, Nazmul MHM<sup>5</sup>

ata, citation and similar papers at [core.ac.uk](https://core.ac.uk)

brought to you by  CORE

provided by UMP Institutional Repository

<sup>2</sup>Jeffrey Cheah School of Medicine and Health Sciences, Monash University, No.8, Jalan Masjid Abu Bakar, 80100 Johor Bahru, Malaysia

<sup>3</sup>Faculty of Science, Lincoln University, 12-18, Jalan SS6/12, Off Jalan Perbandaran, 47301 Petaling Jaya, Selangor Malaysia.

<sup>4</sup>Faculty of Medicine, University Teknologi MARA, Jalan Hospital, Sg Buloh, Selangor 47000, Malaysia

<sup>5</sup>Center of Research Excellence, Graduate School of Medicine, Perdana University, Jalan MAEPS Perdana, Serdang, 43400 Selangor, Malaysia

## Abstract

**Background:** *Pseudomonas aeruginosa* has emerged as a significant opportunistic bacterial pathogen that causes nosocomial infections in healthcare settings resulting in treatment failure throughout the world. This study was carried out to compare the relatedness between virulence characteristics and  $\beta$ -lactamase encoding genes producing *Pseudomonas aeruginosa*.

**Methods:** A total of 120 *P. aeruginosa* isolates were obtained from both paediatric and adult patients of Selayang Hospital, Kuala Lumpur, Malaysia. Phenotypic methods were used to detect various virulence factors (Phospholipase, Hemolysin, Gelatinase, DNase, and Biofilm). All the isolates were evaluated for production of extended spectrum beta-lactamase (ESBL) as well as metallo  $\beta$ -lactamase (MBL) by Double-disk synergy test (DDST) and E-test while AmpC  $\beta$ -lactamase production was detected by disk antagonism test.

**Results:** In this study, 120 *Pseudomonas aeruginosa* isolates (20 each from blood, wounds, respiratory secretions, stools, urine, and sputum samples) were studied. Among *Pseudomonas aeruginosa* isolates, the distribution of virulence factors was positive for hemolysin (48.33%), DNase (43.33%), phospholipase (40.83%), gelatinase (31.66%) production and biofilm formation (34%) respectively. The prevalence of multiple  $\beta$ -lactamase in *P. aeruginosa* showed 19.16% ESBL, 7.5% MBL and 10.83% AmpC production respectively.

**Conclusion:** A regular surveillance is required to reduce public health hazard and the spread of virulence factors and  $\beta$ -lactamase genes among clinical isolates of *Pseudomonas aeruginosa*

**Keywords:** *Pseudomonas aeruginosa*; ESBL; MBL; Virulence factors